


SI Session: Review Exam II
Mondays: 1:30 PM – 3:00 PM & 4:50
PM – 6:20 PM
Wednesdays: 1:30 PM – 3:00 PM &
4:50 PM – 6:20 PM
Room 1239 SNAD(Wed. early rm. 1121)

Prof. Stockton : Calculus III
Fall 2008
SI Leader : Neil Jody

[1] Let $f(x, y) = 4x^2 + 9y^2$.

- (a) Sketch the level curve of f containing the point $(0, -2)$.




- (b) Find the unit vector which points in the direction in which f decreases most rapidly at $(0, -2)$.

[2] Find an equation of the plane tangent to the surface $x^2 + yz^3 = 4$ at the point $(-1, 3, 1)$.

[3] Find the directional derivative of the function $f(x, y, z) = xyz$ at the point $(1, 2, -2)$ in the direction from $(1, 2, -2)$ to $(-1, 0, -1)$.

- [4] Find the absolute extrema of the function $f(x, y) = x^2 + y^2 - 6y$ on the closed region bounded by the graphs $y = 4 - x^2$ and $x + y = 2$.



- [5] Use Lagrange Multipliers to find the maximum and minimum of the function $f(x, y, z) = x + y + z$ on the sphere $x^2 + y^2 + z^2 = 4$.

- [6] Let $f(x, y, z) = x \ln y + y^2 \sin(xz)$. Calculate f_{yxy} .

- [7] Evaluate the integral $\int_1^2 \int_0^{\sqrt{4-x^2}} \frac{x}{x^2 + y^2} dy dx$ by first converting to polar coordinates.



- [8] Let $z = x^2 + xy$, $x = rs + 2t$, $y = r^2 - st$. Calculate z_r when $r = 1$, $s = -2$, and $t = 3$.

[9] Let $f(x, y) = 6x^2 - 2x^3 + 3y^2 + 6xy$. Find all relative extrema and saddle points for f .

- [10] Let D be the triangular region in the xy -plane bounded by the graphs of $x + y = 6$, $y = 2x$, and $5y = x$. Using the change of variables $x = 5u + v$ and $y = u + 2v$, express the integral $\iint_D (5y - x)e^{y-2x} \, dx \, dy$ as an iterated integral in the variables u and v . Do not evaluate the integral.

- [11] Let D be the region in the xy -plane bounded by the x -axis, the y -axis, the line $y = 1$, and the curve $y = \ln x$. Express $\iint_D f(x, y) dA$ as an iterated integral. Do not evaluate the integral.



- [12] Express $\int_0^1 \int_3^{4-x^2} f(x, y) dy dx$ as an iterated integral with the reverse order of integration.



- [13] Let D be the region in the xy -plane bounded by the lines $y = 3x$, $2y = x$, and $x = 4$. Using the change of variables $x = u - 2v$ and $y = 3u - v$, evaluate


$$\iint_D (y - 3x) dy dx .$$



- [14] Let D be the region in the xy -plane bounded by the lines $x + 2y = 2$, $y = x + 1$, and $y = -2x + 4$. Use the change of variables $x = u + 2v$ and $y = u - v + 1$ to evaluate the integral $\iint_D (x - y) dA$.



[15] Evaluate the integral $\int_{-2}^2 \int_0^{\sqrt{4-y^2}} \sin(x^2 + y^2) dx dy$ by first converting to polar coordinates.



- [16] Let D be the region in the xy -plane bounded on the left by the y -axis, above by the graph of $x^2 + y^2 = 4$ and below by the line $y = 1$.

Evaluate $\iint_D \frac{1}{(x^2 + y^2)^{\frac{3}{2}}} dx dy$ by converting to polar coordinates.

